

Invasive competition with Fokker-Planck diffusion and noise

Michael Bengfort^{a,*}, Ivo Siekmann^b, Horst Malchow^a

^a*Institute of Environmental Systems Research, School of Mathematics / Computer Science,
Osnabrück University, Barbarastrasse 12, 49076 Osnabrück, Germany*

^b*Institute for Mathematical Stochastics, Georg-August-University of Göttingen,
Goldschmidtstraße 7, 37077 Göttingen, Germany*

Abstract

Defeat and success of the competitive invasion of a populated area is described with a standard Lotka-Volterra competition model. The resident is adapted to the heterogeneous living conditions, i.e., its motion is modelled as space-dependent, so-called Fokker-Planck diffusion. The invader's diffusion is taken as neutral Fickian. Furthermore, it is studied how multiplicative environmental noise fosters or hinders the invasion.

Keywords: competition, invasion, Fokker-Planck diffusion, environmental noise

1. Introduction

Interactions and movements of populations in a heterogeneous and variable environment are often modelled with stochastic reaction-diffusion equations. Diffusive fluxes in ecology can differ due to specifics of the population's relationships and environmental heterogeneity. They might be neutral cf. eq. (1), attractive (2) or repulsive (3), i.e., for N populations

$$\vec{j}_{in} = -D_i(\vec{r}, \mathbf{X}) \vec{\nabla} X_i(\vec{r}, t), \quad (1)$$

$$\vec{j}_{ia} = -D_i^2(\vec{r}, \mathbf{X}) \vec{\nabla} \left[\frac{X_i(\vec{r}, t)}{D_i(\vec{r}, \mathbf{X})} \right] = +X_i(\vec{r}, t) \vec{\nabla} D_i(\vec{r}, \mathbf{X}) - D_i(\vec{r}, \mathbf{X}) \vec{\nabla} X_i(\vec{r}, t), \quad (2)$$

$$\vec{j}_{ir} = -\vec{\nabla} [D_i(\vec{r}, \mathbf{X}) X_i(\vec{r}, t)] = -X_i(\vec{r}, t) \vec{\nabla} D_i(\vec{r}, \mathbf{X}) - D_i(\vec{r}, \mathbf{X}) \vec{\nabla} X_i(\vec{r}, t); \quad (3)$$

$$i = 1, 2, \dots, N.$$

The usual notation is used: $\mathbf{X}(\vec{r}, t) = \{X_i(\vec{r}, t); i = 1, 2, \dots, N\}$ is the vector of population densities at position $\vec{r} = \{x, y\}$ and time t and $D_i(\vec{r}, \mathbf{X})$ their possibly space- and density-dependent diffusion coefficient. The formulations (1–3) have been elaborated by Skellam

*Corresponding author

Email addresses: michael.bengfort@uni-osnabrueck.de (Michael Bengfort),
ivo.siekmann@mathematik.uni-goettingen.de (Ivo Siekmann), horst.malchow@uni-osnabrueck.de
(Horst Malchow)

(1951; 1973, and nicely summarized by Okubo (1980), see also Aronson (1985) and Murray (1989). In order to complete the list of ecodiffusive fluxes in heterogeneous media, one could add the flux in environmental potentials $U(\vec{r})$

$$\vec{j}_{ip} = \vec{j}_{ik} + \gamma_i X_i(\vec{r}, t) \vec{\nabla} U(\vec{r}); \quad i = 1, 2, \dots, N; \quad (4)$$

where γ_i is called the coefficient of affinity of X_i to the environment and index k can be n , a and r respectively, i.e., one of the fluxes (1–3) can be applied. The minima of $U(\vec{r})$ correspond to preferable and, therefore, attracting habitats. The latter concept has been derived from the ideas of habitat value and environmental density (Morisita, 1971; Shigesada and Teramoto, 1978).

The neutral diffusion is also called Fickian (Fick, 1855) whereas the repulsive type is named after Fokker and Planck (1914; 1917). For a certain density dependence of diffusion, the latter has been used for modelling the spatial segregation of populations (Shigesada et al., 1979; Mimura and Kawasaki, 1980) as well as the formation of Turing patterns (Malchow, 1988).

In a recent publication (Bengfort et al., 2016), the diffusivities have been assumed purely space-dependent. Spatial patterns may already occur without any interactions. For this setting, the spatially stationary solution has been derived. Furthermore, the speed of diffusive waves of a single logistically growing population has been analytically estimated, and conditions for the formation of spatio-temporal and Turing patterns in an excitable prey-predator system have been given.

Another recent publication (Siekman and Malchow, 2016) has dealt with the control of invasion of a populated area by selective infection of the invader as well as by white and coloured noise-modulated environments the resident is adapted to but being unfavourable for the invading population.

The present work shall link the two latter approaches. The Lotka-Volterra textbook model of the competition of two populations is combined with space-dependent Fokker-Planck diffusion of the residents, Fickian diffusion of the invaders and environmental noise. It will be shown that the spatial heterogeneity modelled by Fokker-Planck diffusion but also the external noise can foster or hinder the invasion.

2. The stochastic competition-diffusion model

The dynamics of resident X_1 and invader X_2 is described by

$$\frac{\partial X_1}{\partial t} = (1 - X_1)X_1 - c_{12}X_1X_2 + d_1\nabla^2(X_1D^*(x, y)) + g_1(X_1)\xi(\vec{r}, t), \quad (5)$$

$$\frac{\partial X_2}{\partial t} = (1 - X_2)X_2 - c_{21}X_1X_2 + d_2\nabla^2X_2 + g_2(X_2)\xi(\vec{r}, t). \quad (6)$$

The space dependence of the resident's diffusivity is chosen as

$$D^*(x, y) = D_0 + \begin{cases} a \left(\sin(\sqrt{x^2 + y^2}) \right)^m & \text{if } \sqrt{x^2 + y^2} < 3\pi, \\ a (\sin(3\pi))^m & \text{else.} \end{cases} \quad (7)$$

This spatially varying diffusivity is meant to represent a simple fragmented landscape with a varying habitat quality for species X_1 . The parameter m is an even number which controls the steepness of D^* .

For simplicity, just uncorrelated white noise $\xi(\vec{r}, t)$ is applied here, i.e.,

$$\langle \xi(\vec{r}, t) \rangle = 0, \langle \xi(\vec{r}_1, t_1) \xi(\vec{r}_2, t_2) \rangle = \delta(\vec{r}_1 - \vec{r}_2) \delta(t_1 - t_2) \quad (8)$$

with linearly density dependent noise intensities

$$g_i(X_i) = \omega_i X_i; i = 1, 2. \quad (9)$$

3. Numerical methods

3.1. Crank-Nicolson scheme for two dimensions with Fokker-Planck diffusion

We split the Laplace operator into two parts. First, we calculate the diffusion in one spatial dimension (x), second we do the same for the other spatial dimension (y).

$$\frac{\partial X}{\partial t} = \vec{\nabla}^2(XD) = \frac{\partial^2(XD)}{\partial x^2} + \frac{\partial^2(XD)}{\partial y^2}, \quad (10)$$

where X is the population density and D its spatially varying diffusion coefficient which can be written as

$$D(x, y) = d_1 D^*(x, y) \quad (11)$$

with $d_1 = \text{const}$ and $D^*(x, y) \neq 0 \forall x, y$. Now we formulate the Crank-Nicolson algorithm (Crank and Nicolson, 1947) for one spatial dimension as follows

$$\begin{aligned} \frac{X_k^{t+\Delta t} - X_k^t}{\Delta t} = \frac{d_1}{2\Delta x^2} & \left(X_{k+1}^{t+\Delta t} D_{k+1}^* - 2X_k^{t+\Delta t} D_k^* + X_{k-1}^{t+\Delta t} D_{k-1}^* \right. \\ & \left. + X_{k+1}^t D_{k+1}^* - 2X_k^t D_k^* + X_{k-1}^t D_{k-1}^* \right). \end{aligned} \quad (12)$$

Here $k \in (1, n)$ is the index of the spatial position of X , whereas t is the time which varies with a discrete time step Δt . With $\alpha = d_1 \frac{\Delta t}{\Delta x^2}$ we can write this as a system of linear equations

$$\mathbf{A} \left(\vec{X}^{t+\Delta t} \vec{D}^* \right) = \mathbf{B} \left(\vec{X}^t \vec{D}^* \right) \quad (13)$$

where $\vec{X}$ and $\vec{D}^*$ are vectors of length n including the values of X_k and D_k^* at each spatial position in one dimension $k \in (1, n)$. $\mathbf{A}$ and $\mathbf{B}$ are $n \times n$ tridiagonal matrices

$$\mathbf{A} = \begin{pmatrix} 2 \left(\frac{1}{D_1^*} + \alpha \right) & -\alpha & 0 & \dots & 0 \\ -\alpha & \ddots & -\alpha & 0 & \vdots \\ 0 & -\alpha & \ddots & \ddots & 0 \\ \vdots & \dots & \ddots & \ddots & -\alpha \\ 0 & \dots & & -\alpha & 2 \left(\frac{1}{D_n^*} + \alpha \right) \end{pmatrix}$$

and

$$\mathbf{B} = \begin{pmatrix} 2 \left(\frac{1}{D_1^*} - \alpha \right) & \alpha & 0 & \dots & 0 \\ \alpha & \ddots & \alpha & 0 & \vdots \\ 0 & \alpha & \ddots & \ddots & 0 \\ \vdots & \dots & \ddots & \ddots & \alpha \\ 0 & \dots & & \alpha & 2 \left(\frac{1}{D_n^*} - \alpha \right) \end{pmatrix}.$$

This implicit scheme has been proven to be unconditionally stable for two spatial dimensions.

In order to implement zero-flux boundary conditions we have to add the term $-\alpha$ to the matrix components $\mathbf{A}_{11}$ and $\mathbf{A}_{nn}$, and the term α to the matrix components $\mathbf{B}_{11}$ and $\mathbf{B}_{nn}$. To calculate the distribution of X^t at time step $t + \Delta t$, we have to multiply the vector $\vec{X}^t$ with the spatially varying coefficient of diffusion $\vec{D}^*$ and solve the equation $\mathbf{A}\vec{Y} = \mathbf{B}\vec{X}$, where $\vec{X}$ is a input-vector (in our case $\vec{X}^t \cdot \vec{D}^*$) and $\vec{Y}$ is a output-vector. After that the components of the output-vector $\vec{Y}$ has to be divided with the corresponding components of the vector $\vec{D}^*$, which is temporally constant in order to get the distribution $X^{t+\Delta t}$. Once this scheme has been performed for each row in one spatial direction it has to be repeated for the other spatial dimension in every time step.

3.2. Derivative-free Milstein method for interactions and noise

For numerical integration of the interaction and noise terms, the derivative-free Milstein method is used (Milstein, 1995; Kloeden and Platen, 1999). The Milstein scheme reads for white noise (8,9) with time step Δt and in Stratonovich interpretation

$$X_i^{t+\Delta t} = X_i^t + f_i(X_i^t)\Delta t + \omega_i X_i^t \Delta W_i + \frac{\omega_i}{2} \left[f_i(X_i^t) \sqrt{\Delta t} + \omega_i X_i^t \right] (\Delta W_i)^2, \quad (14)$$

with

$$\Delta W_i = W_i^{t+\Delta t} - W_i^t \sim \sqrt{\Delta t} \mathcal{N}(0, 1).$$

As usual, $\mathcal{N}(0, 1)$ stands for the normal distribution with zero mean and unity variance. The required uniformly distributed random numbers are generated with the Mersenne Twister (Matsumoto and Nishimura, 1998), the normally distributed random numbers with the common Box-Muller algorithm (Box and Muller, 1958).

4. Numerical simulations and results

The following parameters have been applied:

$$D_0 = 1, \quad m = 8, \quad c_{12} = c_{21} = 1.2.$$

Because both species are described with the same parameter values, the difference in the coefficient of diffusion determines whether or not an invasion of species X_2 is successful in case of homogeneous D , i.e., $a = 0$, cf. eq. (7).

If the native species X_1 has a smaller coefficient of diffusion in certain areas of the domain, whereas in the other areas its coefficient of diffusion $d_1 \cdot D^*$ is larger as the constant coefficient d_2 , invasion is successful in those areas where the invader has the larger coefficient of diffusion (Fig. 2). Areas with a high diffusivity of the native species act as barrier for the invasion. This fits well earlier published results on diffusion-controlled competitive invasions (Malchow et al., 2011). In this scenario multiplicative density-dependent noise, as described in eq. (8) and (9), accelerates the speed of invasion (Fig. 2b). Strong noise can push the invader through the barriers of large resident diffusivity and induce invasions in other regions with low resident diffusivity.

Because of the Fokker-Planck diffusion in eq. (5), the spatial distribution of the resident species, X_1 , develops proportional to $\nabla^2 D^*(x, y)$, as described in Bengfort et al. (2016). If this effect is strong enough, the reduced resident concentration in areas with high resident

diffusivity enables an invasion of species X_2 , even if the diffusivity of X_1 is larger than the diffusivity of X_2 everywhere in the domain (Fig. 3). In this scenario, multiplicative density-dependent noise has a decelerating effect on the speed of invasion (Fig. 3b).




Figure 1: Initial settings for densities (green = resident, red = invader) and resident's diffusivity




Figure 2: $d_1 = 5$, $d_2 = 25$, $a = 9$: The density of the resident species is reduced in areas of large D^* . The invader successfully invades the space, where it has a larger coefficient of diffusion as the resident species. Density-dependent multiplicative noise accelerates the invasion in areas of small D^* . Areas with large D^* act as a barrier for the invasion. Strong noise can break through these barriers and induces invasion of X_2 in the inner circles with small D^* .




Figure 3: $t = 4400$, $d_1 = 30$, $d_2 = 25$, $a = 19$; large $\nabla^2 D(\vec{r})$: Due to the reduced resident concentration in areas of large D^* invasion is possible even though the invader has a smaller coefficient of diffusion everywhere in the spatial domain. Noise reduces the invasion speed. Strong noise can invert the invasion.

5. Conclusions

It has been shown that a non-uniform diffusivity, i.e., Fokker-Planck diffusion, of a resident species in a spatially heterogeneous habitat can have different effects on the ability of a similar competing species to invade the habitat.

This Fokker-Planck type of modelling the movement of organisms generates patterns in the spatial population distribution which correspond to the spatial variation of the diffusion coefficient. If this effect is small, the competitor can invade the domain in areas where its (spatially constant) coefficient of diffusion is larger than that of the resident species. This is not surprising because both species are described with equal parameters for growth and competition so that diffusivity determines the success of invasion if the size of the initial patch of the invading species exceeds the related critical patch size. In a non-deterministic environment, where the populations are subject to stochastic fluctuations, the speed of invasion increases with increasing noise intensity. Strong noise can also induce invasions in areas which are perfectly protected against an invasion in the deterministic case. If the pattern forming effect of the Fokker-Planck diffusion is stronger, invasion is possible even though the coefficient of the invader is smaller than the one of the resident species everywhere in the domain. Contrary to the former example, noise has a negative effect on the success of invasion. This is caused by the fact, that the density dependent noise counteract the pattern forming properties of the Fokker-Planck diffusion. The resident species benefits from the homogenising effect of the noise because it has a larger coefficient of diffusion than the invader.

In this paper, Gaussian noise in time and space was applied in order to model the variability of the environment. For future research it would be interesting to investigate the effect of spatially and/or temporally coloured noise in combination with the Fokker-Planck diffusion which generates patterns in the resident species with a certain wavelength.

Here, it was assumed that only the resident species favours certain areas in the domain and consequently move with a spatially varying speed and is therefore described with Fokker-Planck diffusion. One can also think of situations where the invader is described with a heterogeneous coefficient of diffusion as well. The areas favoured by the invading species can be the same as for the resident or independently distributed.

Acknowledgements

The authors acknowledge the stimulating working and living conditions at the Mediterranean Institute of Oceanography during several visits of Aix-Marseille University. They appreciated the professional cooperation but also the sincere hospitality of Jean-Christophe Poggiale and his colleagues. Last but not least, H.M. is thankful for the perfect organization of MPDE'16.

References

- Aronson, D.G., 1985. The role of diffusion in mathematical population biology: Skellam revisited, in: Capasso, V., Grosso, E., Paveri-Fontana, S.L. (Eds.), *Mathematics in Biology and Medicine*. Springer, Berlin. volume 56 of *Lecture Notes in Biomathematics*, pp. 2–6.
- Bengfort, M., Malchow, H., Hilker, F.M., 2016. The Fokker-Planck law of diffusion and pattern formation in heterogeneous media. *Journal of Mathematical Biology* 73, 683–704.
- Box, G.E.P., Muller, M.E., 1958. A note on the generation of random normal deviates. *Annals of Mathematical Statistics* 29, 610–611.
- Crank, J., Nicolson, P., 1947. A practical method for numerical evaluation of solutions of partial differential equations of the heat-conduction type, in: *Mathematical Proceedings of the Cambridge Philosophical Society*, pp. 50–67.
- Fick, A., 1855. Ueber Diffusion. *Annalen der Physik* 170, 59–86 (in German).

- Fokker, A.D., 1914. Die mittlere Energie rotierender elektrischer Dipole im Strahlungsfeld. *Annalen der Physik* 348, 810–820 (in German).
- Kloeden, P.E., Platen, E., 1999. Numerical solution of stochastic differential equations. volume 23 of *Applications of Mathematics*. Springer, Berlin.
- Malchow, H., 1988. Spatial patterning of interacting and dispersing populations. *Memoirs of the Faculty of Science, Kyoto University (Series of Biology)* 13, 83–100.
- Malchow, H., James, A., Brown, R., 2011. Competitive and diffusive invasion in a noisy environment. *Mathematical Medicine and Biology* 28, 153–163.
- Matsumoto, M., Nishimura, T., 1998. Mersenne Twister: a 623-dimensionally equidistributed uniform pseudorandom number generator. *ACM Transactions on Modeling and Computer Simulation* 8, 3–30.
- Milstein, G.N., 1995. Numerical integration of stochastic differential equations. volume 313 of *Mathematics and Its Applications*. Kluwer Academic Publishers, Dordrecht.
- Mimura, M., Kawasaki, K., 1980. Spatial segregation in competitive interaction-diffusion equations. *Journal of Mathematical Biology* 9, 49–64.
- Morisita, M., 1971. Measuring of habitat value by the “environmental density” method, in: Patil, C.D., Pielou, E.C., Waters, W.E. (Eds.), *Spatial patterns and statistical distributions*. The Pennsylvania State University Press, University Park. volume 1 of *Statistical Ecology*, pp. 379–401.
- Murray, J.D., 1989. Mathematical biology. volume 19 of *Biomathematics Texts*. Springer, Berlin.
- Okubo, A., 1980. Diffusion and ecological problems: Mathematical models. volume 10 of *Biomathematics Texts*. Springer, Berlin.
- Planck, M., 1917. Über einen Satz der statistischen Dynamik und seine Erweiterung in der Quantentheorie. *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften* XXIV, 324–341 (in German).
- Shigesada, N., Kawasaki, K., Teramoto, E., 1979. Spatial segregation of interacting species. *Journal of Theoretical Biology* 79, 83–99.

- Shigesada, N., Teramoto, E., 1978. A consideration on the theory of environmental density. Japanese Journal of Ecology 28, 1–8 (in Japanese).
- Siekman, I., Malchow, H., 2016. Fighting enemies and noise: Competition of residents and invaders in a stochastically fluctuating environment. Mathematical Modelling of Natural Phenomena 11, 120–140.
- Skellam, J.G., 1951. Random dispersal in theoretical populations. Biometrika 38, 196–218.
- Skellam, J.G., 1973. The formulation and interpretation of mathematical models of diffusionary processes in population biology, in: Bartlett, M.S., Hiorns, R. (Eds.), The mathematical theory of the dynamics of biological populations. Academic Press, New York, pp. 63–85.